



Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest: composition and floristic relationships with other ombrophilous forests

Samyra Gomes Furtado¹ and Luiz Menini Neto^{1,2,3*}

Received: March 18, 2016

Accepted: July 11, 2016

ABSTRACT

Only a few studies regarding vascular epiphytes have been conducted in mixed ombrophilous forests (MOF) in Serra da Mantiqueira, a mountainous environment in the Brazilian Atlantic Forest, where the relationships of epiphytic flora with other physiognomies are unknown. This study aimed to survey the epiphytes of a MOF remnant located in Serra da Mantiqueira, and to analyze the floristic relationships with ombrophilous forests of the Southern and Southeastern regions of Brazil. The checklist was compared with 51 other areas composed of ombrophilous forests and/or ecotones with other physiognomies using UPGMA (with Sørensen index), and canonical correspondence analysis (CCA). We recorded 138 species, and Orchidaceae and Polypodiaceae were the richest families (51 and 23 species, respectively). The UPGMA showed the importance of physiognomy and elevation in the floristic relationships, and CCA reinforced the influence of elevation, in addition to the shortest distance to the ocean and minimum annual temperature; however, in this analysis, the physiognomies showed little influence on the relationships. The epiphytic flora of MOF of Southern and Southeastern regions of Brazil has different relationships compared with the data available for shrubs and trees, suggesting a greater importance of phorophytic species than geographical distance and, to some extent, environmental variables.

Keywords: biodiversity, conservation, endangered species, environmental variables, epiphytism, mixed ombrophilous forest, Serra da Mantiqueira, similarity

Introduction

Brazilian Atlantic Forest exhibits high diversity, harbouring approximately 16000 plant species, totalling about 46% of the country flora, of which approximately 7500 are endemic (Stehmann *et al.* 2009; Forzza *et al.* 2012). These numbers, together with intense anthropogenic degradation, earned it the status of the world hotspot of biodiversity (Mittermeier *et al.* 2004). However, there are great gaps

in knowledge about the native flora, especially in places of difficult access, such as the mountainous environments (Martinelli 2007; Rapini *et al.* 2009), as well as for some functional groups, such as epiphytes (Kersten 2010).

Mountainous regions are environments with high indices of richness and endemism, in addition to representing islands of vegetation with important forest remnants (Martinelli 2007), maintained due to the barrier represented by the relief, which avoids direct anthropogenic action.

¹ Programa de Pós-graduação em Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Juiz de Fora, 36036-900, Juiz de Fora, MG, Brazil

² Centro de Ensino Superior de Juiz de Fora, 36030-776, Juiz de Fora, MG, Brazil

³ Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Juiz de Fora, 36036-900, Juiz de Fora, Minas Gerais; Brazil

* Corresponding author: furtadosg@gmail.com

The Serra da Mantiqueira covers the borders of Minas Gerais, São Paulo, Rio de Janeiro, and Espírito Santo states, forming together with the Serra do Mar a mountainous range consisting primarily of Atlantic Forest (Rizzini 1997). It is considered a priority for conservation and study due to its biotic and abiotic features (Drummond *et al.* 2005; Lino *et al.* 2007; Saout *et al.* 2013).

Serra do Papagaio is one of the natural areas that must be highlighted in the Serra da Mantiqueira in Minas Gerais. It is protected by a state park (Parque Estadual da Serra do Papagaio [PESP], composed of approximately 23000 ha, and is geographically connected with the Parque Nacional do Itatiaia, representing a continuous montane environment. Despite the importance of this region, only a few floristic/ecologic studies have been conducted to date (Scolforo *et al.* 2008; Pereira *et al.* 2013; Santiago 2013; Furtado & Menini Neto 2015a; Santana 2016).

The PESP harbours one of the rare fragments of mixed ombrophilous forest (MOF) (or araucaria forest) of Minas Gerais (Ab'Saber 2003; Backes 2009), interspersed with “campo de altitude” and dense ombrophilous forest (DOF). This is the only protected MOF fragment by a conservation unity of integral protection in Minas Gerais (Furtado & Menini Neto 2015a). It is one of the most threatened forest ecosystems of the country. It is estimated that only about 3% of the original cover of this physiognomy remains, including exploited and regeneration areas (Bauermann & Behling 2009). This forest formation reaches the highest elevation in the Serra da Mantiqueira (Backes 2009), and, in PESP, this elevation can reach 2000 m a.s.l. (SG Furtado & L Menini Neto unpubl. res.).

Several studies were carried out in Neotropical Region showing the astonishing diversity of vascular epiphytes (*e.g.*, Gentry & Dodson 1987; Catchpole 2004; Benavides *et al.* 2005; Blum *et al.* 2011; Alves & Menini Neto 2014; Leitman *et al.* 2014), as well as the importance of elevation gradient on this diversity, especially in the Andes and Central America (*e.g.*, Krömer *et al.* 2005; Cardelús *et al.* 2006; Watkins Jr. *et al.* 2006; Furtado 2016). However, despite the increasing number of studies on flora and ecology of epiphytes, especially in recent years, there is still a shortage, when considering their ecological importance in the tropical forests (Nadkarni 1984; Nieder *et al.* 2000). Although Brazil has a considerable richness of epiphytes, mainly due to the forest physiognomies of the BAF (Freitas *et al.* 2016; Menini Neto *et al.* 2016), studies of the epiphytic synusia only have been intensive during the past 30 years, mainly concentrating in the Southern Region of the country (Kersten 2010). In order to contribute to the reduction of knowledge gaps regarding this functional group in the Atlantic Forest, the goals of this study were: 1) to evaluate the richness and composition of the vascular epiphytes in the physiognomy of MOF and in the ecotone with DOF in the PESP; 2) to analyse the floristic relationships, and respective influence of environmental variables, between

areas of the Southeastern and Southern regions of Brazil with similar vegetation; 3) to test whether the pattern of floristic relationships of trees and shrubs found in MOF of Serra da Mantiqueira and those of Southern Region is corroborated by epiphytic flora.

Materials and methods

Study area

The PESP is located in the southern region of Minas Gerais in Serra da Mantiqueira (Fig. 1), comprising 22917 ha, between the municipalities of Aiuruoca, Alagoa, Baependi, Itamonte, and Pouso Alto (22.1420S, 44.7328W). The elevations are mainly above 1800 m a.s.l., and the climate is classified as Cwb (according to the Köppen classification), a temperate, highland, tropical climate with dry winters (Silva *et al.* 2008).

The park harbours important remnants of Atlantic Forest, composed of a mosaic of high montane DOF, high montane MOF, and “campo de altitude” (which is a vegetation predominantly composed of open fields with grasses, sometimes with rocky outcrops, also named by Safford (1999) as “Brazilian páramos”). In the studied area, the MOF occurs mainly as fragments of alluvial forest, predominantly on humic and histic cambisols, at elevations ranging from 1600-1700 m a.s.l., along the Santo Agostinho brook (Silva *et al.* 2008). It forms continuous vegetation that is composed of three strata: a canopy of *Araucaria angustifolia* (Bertol.) Kuntze (Araucariaceae) (about 30 m high); a second stratum composed predominantly of *Podocarpus lambertii* Klotsch ex Endl. (Podocarpaceae) (10-15 m high); and a third stratum (up to approximately 8-10 m high) composed of shrubs and treelets of the families Lauraceae, Myrtaceae, Primulaceae, and Winteraceae, among others. This physiognomy exhibits transition areas, with the DOF at 1900-2000 m a.s.l., with few individuals of *A. angustifolia* and near complete absence of *P. lambertii*. *Podocarpus lambertii* also occurs in patches interspersed within the “campo de altitude”, adjacent to the alluvial forest (Furtado & Menini Neto 2015a).

Floristic survey

The floristic survey was conducted through monthly expeditions between April 2012 and September 2013 using the walking method (“método de caminhamento”) (Filgueiras *et al.* 1994) in order to cover the largest possible area of MOF and transition with DOF in each expedition. The fertile specimens were collected, herborised, and deposited in the Herbarium CESJ (acronym according to Thiers [2015]). The plants were photographed in the field and published as a rapid colour guide (Furtado & Menini Neto 2013). The species were classified according to their relationships with




Figure 1. Figure 1. Location of Parque Estadual da Serra do Papagaio (1) and other 52 areas used in multivariate analyses. The numbers of localities are presented in Tab. S1 in supplementary material. ● areas composed by mixed ombrophilous forest from Planalto Meridional (Southern Region), ○ areas composed by mixed ombrophilous forest from Serra da Mantiqueira (Southeastern Region), ▲ areas composed by dense ombrophilous forest from Serra do Mar or Serra da Mantiqueira (Southeastern Region), Δ areas composed by dense ombrophilous forest from Serra do Mar (Southern Region).

phorophytes (Benzing 1990) and identified according to the specialised bibliography, consultation with the herbarium material collection, and specialists.

Evolutionary lineages are according to the APG IV (2016) for angiosperms (eudicotyledons, magnoliids, and monocotyledons) and Christenhusz *et al.* (2011) for the ferns (lycophytes and monilophytes). Orchidaceae genera *Maxillaria*, *Oncidium*, and *Pleurothallis* were considered in a broad sense due to the lack of consensus regarding their delimitations and due to several recent proposals of segregation in several smaller genera.

Multivariate analyses

The composition of vascular epiphytes of the PESP was compared to areas with available lists of vascular epiphytes and some areas with extensive vascular flora surveys that discriminated each life form. We used 52 areas of MOF or DOF, and, in some cases, ecotones with other physiognomies occurring in Southern and Southeastern regions of Brazil, in addition to areas of the Serra da Mantiqueira with elevations similar to the PESP (Tab. S1 in supplementary material). The

data were obtained from published studies and the database of herbaria collections available at the site Specieslink of “Centro de Referência em Informação Ambiental” (CRIA) (<http://www.splink.org.br>). All unidentified species were excluded, resulting in a matrix of presence (1) and absence (0), with 910 species.

The similarity between the aforementioned areas was evaluated through cluster analysis using the unweighted pair-group method with arithmetic mean (UPGMA) and similarity index of Sørensen. The cophenetic coefficient was calculated to test the fit between the matrix and resulting dendrogram. A Mantel test was conducted to evaluate the correlation between the geographic distance and calculated similarity among the areas. These analyses were conducted using the software PAST v. 3.01 (Hammer *et al.* 2001).

In order to evaluate the correlation among the environmental variables and composition of vascular epiphytes, a canonical correspondence analysis (CCA) was conducted (ter Braak 1986; Palmer 1993). Previous analyses were performed with a set of 19 climatic variables (annual mean temperature, mean monthly temperature range, isothermality, temperature seasonality, max temperature

Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest: composition and floristic relationships with other ombrophilous forests

of warmest month, min temperature of coldest month, temperature annual range, mean temperature of wettest quarter, mean temperature of driest quarter, mean temperature of warmest quarter, mean temperature of coldest quarter, annual precipitation, precipitation of wettest month, precipitation of driest month, precipitation seasonality, precipitation of wettest quarter, precipitation of driest quarter, precipitation of warmest quarter, precipitation of coldest quarter). The variables were presented by Hijmans *et al.* (2005), and are available in the site WorldClim (www.worldclim.org). These variables were complemented with other considered important to the epiphytic flora (Benzing 1990): minimal and maximal elevations and minimal and maximal annual means of temperature obtained in the respective articles, and the shortest distance to the Atlantic Ocean, which was calculated for each area using the software DIVA-GIS v. 7.5 (Hijmans *et al.* 2001), which is used to represent a seasonality gradient.

After this preliminary analysis, the redundant variables, with high values of inflation, were discarded (ter Braak 1986). Three variables resulted as the most representative and correlated with the two first ordination axes: shortest distance to the ocean, elevation, and minimal annual temperature. The permutation test of Monte Carlo was conducted a posteriori in order to evaluate the significance of the canonical correlations at a significance level of 95%

($p < 0.05$) (ter Braak 1986; Palmer 1993). These analyses were conducted with the software CANOCO v. 4.5 (ter Braak & Smilauer 2002).

Results

Floristic survey

We recorded 25 families, 66 genera, and 138 species in the studied area. The ferns were represented by 43 species (31.16% of total) and the angiosperms by 95 species (68.84%). Orchidaceae was the richest family (51 species, 37%), followed by Polypodiaceae (23 species, 17%), Bromeliaceae, and Piperaceae (10 species, 7% each). *Pleurothallis* s.l. was the richest genus with 11 species, followed by *Peperomia*, with 10 species (Tab. 1). The majority of species occurred in the MOF (122 species), of which 62 were exclusive, and 60 were shared with the ecotone with the DOF, with 16 being exclusive to the ecotone.

The richest evolutionary lineages were the monocotyledons (63 species) and the monilophytes (39 species), being Orchidaceae and Polypodiaceae the richest families, respectively. Characteristic holoepiphytes were the most well represented ecological category (107 species), but the number of accidental holoepiphytes families and species, 10 (40% of total) and 19 (14% of total), respectively,

Table 1. List of vascular epiphytes recorded in the Parque Estadual da Serra do Papagaio, Minas Gerais, Brazil.

Families and species	EC	Habitat	Voucher
Ferns			
Licophytes – Vinícius A.O. Dittrich (CESJ)			
Lycopodiaceae (1/4)			
<i>Phlegmariurus acerosus</i> (Sw.) B.Øllg.	CHL	F	Furtado 74
<i>Phlegmariurus biformis</i> (Hook.) B.Øllg.	CHL	F	Furtado 75
<i>Phlegmariurus fontinaloides</i> (Spring) B.Øllg.	CHL	F/E	Furtado 114
<i>Phlegmariurus quadrifarius</i> (Bory) B.Øllg.	CHL	F	Furtado 66
Monilophytes – Vinícius A.O. Dittrich, Filipe S. Souza (CESJ), A. Salino (BHCB)			
Anemiaceae (1/1)	AHL	F	Furtado 88
<i>Anemia phyllitidis</i> (L.) Sw.			
Aspleniaceae (1/5)			
<i>Asplenium aff. inaequilaterale</i> Willd.	AHL	F	Furtado 221
<i>Asplenium auriculatum</i> Sw.	CHL	F/E	Furtado 46
<i>Asplenium auritum</i> Sw.	CHL	F/E	Furtado 37
<i>Asplenium incurvatum</i> Fee	CHL	F/E	Furtado 93
<i>Asplenium serra</i> Langsd. & Fisch.	CHL	E	Souza 960
Dryopteridaceae (2/3)			
<i>Elaphoglossum gayanum</i> (Pérez) T.Moore	CHL	F/E	Furtado 183
<i>Elaphoglossum vagans</i> (Mett.) Hieron.	FHL	F/E	Furtado 129
<i>Rumohra adiantiformis</i> (G.Forst.) Ching	CHL	F/E	Furtado 113
Hymenophylaceae (3/4)			
<i>Hymenophyllum polyanthos</i> (Sw.) Sw.	CHL	F/E	Furtado 64
<i>Polyphlebium angustum</i> (Carmich.) Ebihara & Dubuisson	CHL	F	Furtado 97
<i>Trichomanes anadromum</i> Rosenst	CHL	F	Furtado 276
<i>Trichomanes polypodioides</i> Raddi	CHL	F	Furtado 90
Ophioglossaceae (1/1)			
<i>Ophioglossum palmatum</i> L.	CHL	E	Furtado 267



Table 1. Cont.

Families and species	EC	Habitat	Voucher
Polypodiaceae (12/23)			
<i>Campyloneurum aglaolepis</i> (Alston) de la Sota	CHL	F	Furtado 94
<i>Campyloneurum angustifolium</i> (Sw.) Féé	CHL	F/E	Menini Neto 965
<i>Campyloneurum nitidum</i> (Kaulf.) C.Presl	CHL	F	Furtado 236
<i>Campyloneurum</i> sp.	CHL	F/E	Furtado 216
<i>Ceradenia albidula</i> (Baker) L.E.Bishop	CHL	E	Furtado 132
<i>Cochlidium punctatum</i> (Raddi) L.E.Bishop	CHL	F/E	Furtado 137
<i>Lellingeria aff. pumila</i> Labiak	CHL	E	Furtado 198
<i>Lellingeria apiculata</i> (Kunze ex Klotzsch) A.R.Sm. & R.C.Moran	CHL	F/E	Furtado 110
<i>Leucotrichum organense</i> (Gardner) Labiak	CHL	F/E	Furtado 5
<i>Leucotrichum</i> sp.	CHL	F	Furtado 44
<i>Melpomene flabelliformis</i> (Poir.) A.R.Sm. & R.C.Moran	CHL	F/E	Furtado 198
<i>Melpomene pilosissima</i> (M.Martens & Galeotti) A.R.Sm. & R.C.Moran	CHL	F/E	Furtado 127
<i>Microgramma percussa</i> (Cav.) de la Sota	CHL	F	Salimena 2836
<i>Microgramma squamulosa</i> (Kaulf.) de la Sota	CHL	F/E	Furtado 8
<i>Micropolypodium achilleifolium</i> Labiak & F.B.Matos	CHL	F	Furtado 150
<i>Pecluma pectinatiformis</i> (Lindm.) M.G.Price	CHL	F/E	Souza 955
<i>Pecluma</i> sp.	CHL	F	Furtado 7
<i>Pleopeltis hirsutissima</i> (Raddi) de la Sota	CHL	F/E	Furtado 32
<i>Pleopeltis macrocarpa</i> (Bory ex Willd.) Kaulf.	CHL	F/E	Furtado 9
<i>Pleopeltis pleopeltidis</i> (Fée) de la Sota	CHL	F/E	Furtado 116
<i>Serpocaulon catharinae</i> (Langsd. & Fisch.) A.R.Sm.	CHL	F/E	Furtado 50
<i>Zygophlebia longipilosa</i> (C.Chr.) L.E.Bishop	CHL	F	Souza 1006
Polypodiaceae indet.	CHL	F	Furtado 45
Pteridaceae (1/2)			
<i>Vittaria graminifolia</i> Kaulf.	CHL	F/E	Souza 979
<i>Vittaria lineata</i> (L.) SM	CHL	F/E	Furtado 136
Angiosperms			
Magnoliids			
Piperaceae (1/10) – Daniele Monteiro (RB)			
<i>Peperomia campinasana</i> C.DC.	CHL	F	Furtado 73
<i>Peperomia catharinae</i> Miq.	CHL	F	Furtado 106
<i>Peperomia cf. glabella</i> (Sw.) A.Dietr.	CHL	F	Furtado 202
<i>Peperomia hilariana</i> Miq.	FHL	F/E	Menini Neto 814
<i>Peperomia hispidula</i> (Sw.) A. Dietr.	CHL	F	Furtado 72
<i>Peperomia mandiocana</i> Miq.	CHL	F/E	Furtado 99
<i>Peperomia subternifolia</i> Yunck.	CHL	F/E	Furtado 98
<i>Peperomia tetraphylla</i> (G.Forst.) Hook. & Arn.	CHL	F/E	Furtado 195
<i>Peperomia trineura</i> Miq.	CHL	F/E	Menini Neto 841
<i>Peperomia trineuroides</i> Dahlst.	CHL	F/E	Furtado 172
Monocotyledons			
Araceae (1/1)			
<i>Philodendron</i> sp.	HEM	E	Nardy 2
Bromeliaceae (5/10) – Rafaela C. Forzza (RB)			
<i>Aechmea aiuruocensis</i> Leme	FHL	E	Furtado 259
<i>Aechmea distichantha</i> Lem.	FHL	F	Furtado 234
<i>Billbergia distachia</i> (Vell.) Mez	CHL	F/E	Furtado 91
<i>Nidularium marigoi</i> Leme	FHL	F/E	Furtado 228
<i>Tillandsia recurvata</i> (L.) L.	CHL	F	Furtado 222
<i>Tillandsia stricta</i> Sol.	CHL	F/E	Furtado 100
<i>Tillandsia tenuifolia</i> L.	CHL	F	Furtado 67
<i>Vriesea bituminosa</i> Wawra	FHL	F/E	Furtado 250
<i>Vriesea gigantea</i> Gaudich.	FHL	F/E	Furtado 146
<i>Vriesea sceptrum</i> Mez	FHL	F/E	Menini Neto 792



**Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest:
composition and floristic relationships with other ombrophilous forests**

Table 1. Cont.

Families and species	EC	Habitat	Voucher
Orchidaceae (18/51) – Luiz Menini Neto, Samyra G. Furtado, Camila Nardy (CESJ)			
<i>Bifrenaria stefanae</i> V.P.Castro	CHL	E	Furtado 124
<i>Bulbophyllum granulosum</i> Barb.Rodr.	CHL	F	Furtado 207
<i>Bulbophyllum regnellii</i> Rchb.f.	CHL	F	Furtado 242
<i>Capanemia adelaidae</i> Brade	CHL	F	Furtado 53
<i>Cryptophoranthus jordanensis</i> Brade	CHL	E	Furtado 274
<i>Dryadella lilliputiana</i> (Cogn.) Luer	CHL	F/E	Furtado 141
<i>Encyclia patens</i> Hook.	CHL	F	Furtado 275
<i>Epidendrum chlorinum</i> Barb.Rodr.	CHL	F/E	Furtado 214
<i>Epidendrum mantiqueiratum</i> Porto & Brade	CHL	F/E	Furtado 85
<i>Gomesa gomezooides</i> (Barb.Rodr.) Pabst	CHL	F/E	Furtado 210
<i>Grobya amherstiae</i> Lindl.	CHL	E	Furtado 255
<i>Hadrolaelia coccinea</i> (Lindl.) Chiron & V.P.Castro	CHL	F/E	Furtado 79
<i>Hadrolaelia mantiqueirae</i> (Fowlie) Fowlie	CHL	F/E	Furtado 80
<i>Hadrolaelia pygmaea</i> (Pabst) Chiron & V.P.Castro	CHL	F	Furtado 208
<i>Hapalorchis lineatus</i> (Lindl.) Schltr.	AHL	E	Furtado 252
<i>Hapalorchis micranthus</i> (Barb.Rodr.) Hoehne	AHL	F/E	Furtado 69
<i>Lankesterella gnoma</i> (Kraenzl.) Hoehne	CHL	F/E	Furtado 169
<i>Loefgrenianthus blanche-amesii</i> (Loefgr.) Hoehne	CHL	F/E	Furtado 176
<i>Maxillaria neuwiedii</i> Rchb.f.	CHL	F	Menini Neto 1108
<i>Maxillaria notyliglossa</i> Rchb.f.	CHL	E	Furtado 243
<i>Maxillaria paranaensis</i> Barb.Rodr.	CHL	F	Furtado 36
<i>Maxillaria picta</i> Hook.	CHL	F/E	Furtado 174
<i>Octomeria crassifolia</i> Lindl.	CHL	F/E	Furtado 35
<i>Octomeria geraensis</i> Barb.Rodr.	CHL	F/E	Furtado 277
<i>Octomeria ochroleuca</i> Barb.Rodr.	CHL	F	Furtado 25
<i>Octomeria wawrae</i> Rchb.f.	CHL	F	Furtado 92
<i>Octomeria</i> sp1	CHL	F	Furtado 167
<i>Octomeria</i> sp2	CHL	F	Furtado 168
<i>Octomeria</i> sp3	CHL	F	Furtado 213
<i>Oncidium cogniauxianum</i> Schltr.	CHL	F	Furtado 203
<i>Oncidium divaricatum</i> Lindl.	CHL	F	Furtado 220
<i>Oncidium forbesii</i> Hook.	CHL	E	Menini Neto 776
<i>Oncidium gardneri</i> Lindl.	CHL	F/E	Furtado 215
<i>Oncidium hookeri</i> Rolfe	CHL	F/E	Furtado 241
<i>Oncidium longicornu</i> Mutel	CHL	F/E	Furtado 162
<i>Phymatidium mellobarretoi</i> Hoehne & Williams	CHL	F/E	Furtado 26
<i>Pleurothallis adenochila</i> Loefgr.	CHL	E	Furtado 119
<i>Pleurothallis bocainensis</i> Porto & Brade	CHL	F/E	Furtado 68
<i>Pleurothallis cf. corticicola</i> Schltr. ex Hoehne	CHL	F	Furtado 251
<i>Pleurothallis grobyi</i> Bateman ex Lindl.	CHL	F/E	Furtado 117
<i>Pleurothallis linearifolia</i> Cogn.	CHL	F	Furtado 163
<i>Pleurothallis pleurothalloides</i> (Cogn.) Handro	CHL	F	Furtado 1
<i>Pleurothallis pterophora</i> Cogn.	CHL	E	Furtado 226
<i>Pleurothallis radialis</i> Porto & Brade	CHL	E	Furtado 253
<i>Pleurothallis rostellata</i> Barb.Rodr.	CHL	F	Furtado 161
<i>Pleurothallis rubens</i> Lindl.	CHL	F/E	Furtado 87
<i>Pleurothallis uniflora</i> Lindl.	CHL	F	Menini Neto 1068
<i>Stelis intermedia</i> Poepp. & Endl.	CHL	F	Menini Neto 1059
<i>Stelis papaverensis</i> Rchb.f.	CHL	F/E	Furtado 186
<i>Stelis</i> sp1	CHL	F/E	Furtado 196
<i>Stelis</i> sp2	CHL	F	Furtado 273
Eudicotyledons			
Araliaceae (1/1)			
<i>Hydrocotyle cf. bonariensis</i> Lam.	AHL	F	Furtado 22



Table 1. Cont.

Families and species	EC	Habitat	Voucher
Asteraceae (2/6)			
<i>Ageratum fastigiatum</i> (Gardner) R.M.King & H.Rob.	AHL	F	Furtado 247
<i>Baccharis crispa</i> Spreng.	AHL	F	Furtado 237
Asteraceae sp1	AHL	F	Furtado 268
Asteraceae sp2	AHL	F	Furtado 269
Asteraceae sp3	AHL	F	Furtado 270
Asteraceae sp4	AHL	F	Furtado 271
Cactaceae (1/2) – Diego R. Gonzaga (CESJ)			
<i>Rhipsalis floccosa</i> Salm-Dyck ex Pfeiff.	CHL	F/E	Furtado 107
<i>Rhipsalis pulchra</i> Loefgr.	CHL	F/E	Furtado 139
Caryophyllaceae (1/1)			
<i>Arenaria lanuginosa</i> (Michx.) Rohrb.	AHL	F	Furtado 23
Ericaceae (1/1) – Andressa Cabral (CESJ)			
<i>Agarista oleifolia</i> (Cham.) G.Don	AHL	F	Santiago 604
Gesneriaceae (2/3) – Luciana C. Pereira (CESJ)			
<i>Nematanthus fornix</i> (Vell.) Chautems	FHL	E	Furtado 160
<i>Sinningia cooperi</i> (Paxton) Wiehler	CHL	F	Furtado 49
<i>Sinningia douglasii</i> (Lindl.) Chautems	CHL	F	Furtado 164
Melastomataceae (3/3) – Luciana L. Justino (CESJ)			
<i>Leandra carassana</i> (DC.) Cogn.	AHL	F	Furtado 248
<i>Miconia hyemalis</i> A.St.-Hil. & Naudin	AHL	F	Furtado 246
<i>Pleiochiton blepharodes</i> (DC.) Reginato et al.	CHL	F	Furtado 201
Onagraceae (1/1)			
<i>Fuchsia regia</i> (Vell.) Munz	FHL	F	Furtado 165
Plantaginaceae (1/1)			
<i>Plantago</i> sp.	AHL	F	Furtado 272
Poaceae (1/1)			
<i>Chusquea</i> sp.	AHL	F/E	Furtado 254
Polygalaceae (1/1)			
<i>Polygala lancifolia</i> A.St.-Hil. & Moq.	AHL	F	Furtado 65
Ranunculaceae (1/1)			
<i>Anemone sellowii</i> Pritz.	AHL	F	Furtado 89
Solanaceae (1/1)			
<i>Dysochroma viridiflorum</i> (Sims) Miers	HEM	F/E	Furtado 244

Numbers between parentheses after the families names represent the number of genera and species, respectively. Names after families represent the specialists that collaborate in the taxa identification. EC: Ecological categories – HEM: hemiepiphyte; AHL: accidental holoepiphyte; CHL: characteristic holoepiphyte; FHL facultative holoepiphyte. Habitat – F: Mixed Ombrophilous Forest; E: ecotone.

must be highlighted, with Asteraceae being the richest family with six species. Also, Melastomataceae must be noted, exhibiting one characteristic holoepiphyte and two accidental holoepiphytes species (Tab. 2).

Multivariate analyses

The cluster analysis resulted in the dendrogram presented in the Fig. 2, which obtained a cophenetic coefficient of 0.86, showing little distortion between the matrix and graphic. The Mantel test resulted in a positive correlation between the geographic distance and similarity matrix ($r = 0.55$, $p = 0.0001$).

Two main clusters were formed (1 and 2). The first (1) is composed mainly of DOF and shows a division in two other groups, with areas of Serra do Mar and Serra da Mantiqueira at elevations ranging from 500-2879 m a.s.l. in a group (\blacktriangle) and areas of Southern and Southeastern regions at elevations ranging from 0-1000 m a.s.l. in the other group (Δ). Cluster 2 grouped together the MOF but shows a segregation of the southern areas at elevations ranging from 340-1200 m a.s.l. (\bullet) from those of Serra da Mantiqueira (including the PESP) at elevations ranging from 1000-2010 m a.s.l. (\circ).

Despite the existence of a branch composed only of MOF areas, the similarity can be considered low among

**Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest:
composition and floristic relationships with other ombrophilous forests**

Table 2. Evolutionary lineages and respective ecological categories.

Evolutionary lineages and families	Ecological categories							
	CHL (107/78%)		FHL (10/7%)		AHL (19/14%)		HEM (2/1%)	
	N	%	N	%	N	%	N	%
Licophytes (4)								
Lycopodiaceae (4)	4	100%	-	-	-	-	-	-
Monilophytes (39)								
Anemiaceae (1)*	-	-	-	-	1	100%	-	-
Aspleniaceae (5)	4	80%	-	-	1	20%	-	-
Dryopteridaceae (3)	2	66.6%	1	33.3%	-	-	-	-
Hymenophyllaceae (4)	4	100%	-	-	-	-	-	-
Ophioglossaceae (1)	1	100%	-	-	-	-	-	-
Polypodiaceae (23)	23	100%	-	-	-	-	-	-
Monilophytes (39)								
Pteridaceae (2)	2	100%	-	-	-	-	-	-
Magnoliids (10)								
Piperaceae (10)	9	90%	1	10%	-	-	-	-
Monocotyledons (63)								
Araceae (1)	-	-	-	-	-	-	1	100%
Bromeliaceae (10)	4	40%	6	60%	-	-	-	-
Orchidaceae (51)	49	96.1 %	-	-	2	3.%	-	-
Poaceae (1)*	-	-	-	-	1	100%	-	-
Eudicotyledons (22)								
Araliaceae (1)*	-	-	-	-	1	100%	-	-
Asteraceae (6)*	-	-	-	-	6	100%	-	-
Cactaceae (2)	2	100%	-	-	-	-	-	-
Caryophyllaceae (1)*	-	-	-	-	1	100%	-	-
Ericaceae (1)*	-	-	-	-	1	100%	-	-
Gesneriaceae (3)	2	66.6%	1	33.3%	-	-	-	-
Melastomataceae (3)	1	33.3%	-	-	2	66.6%	-	-
Onagraceae (1)	-	-	1	100%	-	-	-	-
Plantaginaceae (1)*	-	-	-	-	1	100%	-	-
Polygalaceae (1)*	-	-	-	-	1	100%	-	-
Ranunculaceae (1)*	-	-	-	-	1	100%	-	-
Solanaceae (1)	-	-	-	-	-	-	1	100%

The number of species of each family or evolutionary lineage is between parentheses after their names. CHL – characteristic holoepiphytes; FHL – facultative holoepiphytes; AHL – accidental holoepiphytes; HEM – hemiepiphytes. N – number of species, % – percentage of species of each family distributed in the ecological categories.

* Families with species typically terricolous and represented in this study only by accidental holoepiphytes.

the two aforementioned subsets (around 0.25). Even the subset composed only of the remnants of MOF in the Serra da Mantiqueira exhibited low similarity and shared only 11 species (Fig. 3).

The results of the CCA, highlighted in the Tab. 3, showed eigenvalues higher than 0.3, which is considered high according to Felfili *et al.* (2011), representing a strong gradient in both axes. The values of species-environment correlations also are considered high (0.985 and 0.947 for axes 1 and 2, respectively). The Monte Carlo test showed a significant correlation between the distribution of species and the environmental variables used in the analysis ($p < 0.05$) (Tab. 3). The variables elevation and minimum annual

temperature showed higher correlations with axis 1, while the shortest distance to the ocean was more correlated with axis 2 (Tab. 4).

The ordination diagram (Fig. 4) did not show a clear group among the areas with same physiognomy, as presented in the dendrogram (Fig. 2), especially regarding the MOF; only a tendency of grouping among them was observed. However, the area surveyed in the present study, ‘mgpesp’, was closely related to at least one of the MOF areas in the Serra da Mantiqueira, ‘sppec’ (Parque Estadual de Campos do Jordão, in São Paulo state), and both were more correlated with the areas of DOF at high elevations than those of MOF occurring in the Southern Region of Brazil (Fig. 4).




Figure 2. Dendrogram (Sørensen similarity index) obtained in the similarity analysis with 53 localities of the Southeastern and Southern regions of Brazil based on a binary matrix of 910 species of vascular epiphytes. Cophenetic coefficient = 0.86. Numbers in the branches are explained in the text. DOF: dense ombrophilous forest; MOF: mixed ombrophilous forest; CR: 'campo rupestre'; CA: 'campo de altitude'; SSF: seasonal semi-deciduous forest; RES: 'restinga' (coastal vegetation); MAN: Mangrove. ● areas composed by mixed ombrophilous forest from Planalto Meridional (Southern Region), ○ areas composed by mixed ombrophilous forest from Serra da Mantiqueira (Southeastern Region), ▲ areas composed by dense ombrophilous forest from Serra do Mar or Serra da Mantiqueira (Southeastern Region), Δ areas composed by dense ombrophilous forest from Serra do Mar (Southern Region).

Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest:
composition and floristic relationships with other ombrophilous forests




Figure 3. Venn diagram with the superposition of vascular epiphytic species of areas with mixed ombrophilous forest (MOF) of Serra da Mantiqueira: Parque Estadual da Serra do Papagaio; Serra da Pedra Branca; and Parque Estadual de Campos do Jordão. SI: Similarity index of Sørensen.




Figure 4. Bi-plot diagram results of the canonical correspondence analysis showing the relationships of 53 areas of the Southeastern and Southern regions of Brazil based on a binary matrix of 910 species of vascular epiphytes and the main environmental variables. The diagram shows the ordination of the first two axes. ● areas composed by mixed ombrophilous forest from Planalto Meridional (Southern Region), ○ areas composed by mixed ombrophilous forest from Serra da Mantiqueira (Southeastern Region), ▲ areas composed by dense ombrophilous forest from Serra do Mar or Serra da Mantiqueira (Southeastern Region), △ areas composed by dense ombrophilous forest from Serra do Mar (Southern Region).

Table 3. Estimators of the two first axes of canonical ordination of vascular epiphytes of 53 areas of Southeastern and Southern regions as well as the most important environmental variables.

Estimators	Axis 1	Axis 2
Eigenvalues	0.448	0.373
Species-environment correlations	0.985	0.947
Cumulative percentage variance of species-data	5.3	9.8
Cumulative percentage variance of species-environment relation	42.2	77.4
Monte Carlo test (p)	0.002	0.002*

*All canonical axes.

Table 4. Correlations of the environmental variables with the two first axes of canonical ordination of vascular epiphytes of 53 areas of Southeastern and Southern regions.

Environmental variables	Correlation coefficients	
	Axis 1	Axis 2
Shortest distance to the ocean	0.2431	0.9098
Elevation	0.9626	-0.1219
Minimum annual temperature	-0.3739	-0.0116

Discussion

Floristic survey

The species richness of each evolutionary lineage is similar to that observed in the Atlantic Forest (Kersten 2010; Freitas *et al.* 2016), although the proportion of representation of each is different. We found a lower percentage of monocotyledons (approximately 46% in the PESP versus approximately 64% for the Atlantic Forest) and a higher percentage of monilophytes (approximately 29% in the PESP versus approximately 16% for the Atlantic Forest). This lower representation of monocotyledons is due to the reduced number of species of Bromeliaceae (10 species) and Araceae (only one species). The monilophytes exhibited a larger contribution to the species composition, as the group is recognisably rich in the MOF, especially due to the Polypodiaceae, according to Kersten (2010).

Despite Orchidaceae, Polypodiaceae, Bromeliaceae, Araceae and Piperaceae are the five richest families in epiphytes in Atlantic domain (Kersten 2010; Freitas *et al.* 2016) and Neotropical region (Gentry & Dodson 1987) as well as at the global level (Zotz 2013), the contribution of each family in this study was different. When considering the physiognomy of MOF only, the first three families also are the richest. However, the Piperaceae contribution increases, being the fourth richest, and the Araceae contribution considerably decreases, according to Kersten (2010). In the present study, this fact is corroborated, since Araceae is represented only by *Philodendron* sp., found in the ecotone between MOF and DOF.

The two richest genera (*Pleurothallis* s.l. and *Peperomia*) are prominent in the Atlantic domain as well as in the Neotropical region. *Pleurothallis* s.l. is one of the richest genera of Orchidaceae and the largest among the epiphytic plants, and the Brazilian Atlantic Forest is one of the centres

of diversity especially in areas of high elevations (Pridgeon 1982; Luer 1986; Gentry & Dodson 1987), such as the PESP. *Peperomia* is one of the largest genera of Piperaceae and exhibits high richness in the Brazilian Atlantic Forest (Menini Neto *et al.* 2016), especially in the ombrophilous forest (Carvalho-Silva 2008), and it is the richest genus among epiphytes if the large genera of Orchidaceae are excluded (Zotz 2013), justifying the number of species.

The richness of vascular epiphytes of the PESP is higher than that of other studied areas in the MOF of the Southern Region (e.g., Cervi & Dombrowski 1985; Cervi *et al.* 1988; Dittrich *et al.* 1999; Kersten & Silva 2002; Borgo & Silva 2003; Kersten 2006; Kersten *et al.* 2009) and in the MOF of Parque Estadual de Campos do Jordão (located at the Serra da Mantiqueira) (Mania 2013), even if we consider only the richness found in the araucaria forests of the PESP (122 species). The PESP exhibits higher richness than that found in several studies conducted in seasonal semi-deciduous forest (Aguiar *et al.* 1981; Dislich & Mantovani 1998; Borgo *et al.* 2002; Rogalski & Zanin 2003; Giongo & Waechter 2004). This contradicts the data gathered by Kersten (2010), who found this physiognomy richer compared to the MOF of Southern Region of Brazil, although this author suggests that status of conservation of the MOF could be the responsible for this result. Thus, the degree of conservation of the PESP must be responsible, in part, for these results, although other features, such as elevation, could influence observed richness (Furtado & Menini Neto 2015a), once a richness peak for epiphytes in altitudinal gradients is common among 1000-2000 m (Madison 1977; Gentry e Dodson 1987; Benzing 1990; Krömer *et al.* 2005; Cardelús *et al.* 2006).

Two other aspects can also influence the richness and must be addressed. The lower latitude of PESP compared with other areas composed by MOF, also can be important due to the influence of latitude on the temperature, which is a relevant feature regarding the epiphyte richness (Benzing

1990). Montane environment itself is another possible influence on the richness found in the present study. Such environment is often found to be a refuge to species and, consequently, shows remarkable richness and endemism if compared with lowland vegetation (Körner 2004; Martinelli 2007).

On the other hand, the ecotone between the MOF and DOF exhibited lower richness (76 species) when compared to the study conducted by Kersten (2006) in a similar environment of transition, in which 143 species were recorded. The same situation occurs when comparing the PESP with areas of DOF, which are typically richer (Breier 2005; Petean 2009; Bonnet *et al.* 2013a, b). However, in some cases, the PESP exhibits a higher richness (Hertel 1950; Petean 2002). Such result must be related to the absence of *Podocarpus lambertii* in the ecotone area. This tree species represents an important phorophyte in the PESP, harbouring 89 of the epiphyte species or 75% of the total recorded in this study (Furtado & Menini Neto 2015a).

Results confirm that characteristic holoepiphyte is the most common ecological category, corroborating similar studies conducted in MOF (Dittrich *et al.* 1999; Hefler & Faustioni 2004; Buzatto *et al.* 2008; Bonnet *et al.* 2011). However, accidental holoepiphytes, as the second-most representative category, is unusual (Bonnet *et al.* 2011) and must be noted. In the PESP, the majority of accidental holoepiphytes was found in some parts of the forest that suffered from fire in the year 2011 that had their entire or almost entire epiphytic communities destroyed.

Anthropogenic disturbances (as fire) are often responsible to alter the community composition, opening space to the establishment of opportunist and/or ruderal species that tolerate the new disturbed environment (Hobbs *et al.* 1992) and occupy the earliest stages of succession (Monaco *et al.* 2002). Thus, such disturbances can be related with the establishment of accidental holoepiphytes in the studied site, consequently enhancing their proportion in comparison with characteristic holoepiphytes. Some weed/ruderal species were already recorded as accidental holoepiphytes in disturbed environments (*e.g.*, *Ageratum conyzoides*, *Drymaria cordata*, *Erechtites valerianaefolius*, *Plantago major*, *Setaria palmifolia*) (Holzner & Numata 1982). Species of these genera were also found as accidental holoepiphytes in the present study (*Ageratum* and *Plantago*) and in some other studies dealing with vascular epiphytes in disturbed environment (for example, Bhatt *et al.* 2015; Furtado & Menini Neto 2015b). It is necessary to conduct more accurate studies in addition to better sampling of this ecological category of epiphytes, which is neglected in several studies regarding epiphyte synusia, and deserves more attention as pointed out by Zott (2013). Moreover, Benzing (1990) emphasised that environments with high moisture facilitate the occurrence of accidental species, which can explain, in part, the representativeness of this category in the present study.

Multivariate analyses

Importance of vegetation formation and elevation in the composition and distribution of vascular epiphytes showed in the dendrogram is similar to the pattern found for angiosperm epiphytes by Menini Neto *et al.* (2009) although these authors used fewer areas than the present study. In the graphic, there is a tendency of grouping the areas that share DOF but segregation of the MOF of Southern and Southeastern regions.

The scatter plot of the CCA reinforced the influence of elevation but added the shortest distance to the ocean and the minimum annual temperature as important in calculation of the relationships. The set of variables of this study were also showed to be relevant in studies dealing with biogeography and floristic relationships of angiosperm epiphytes in Atlantic Forest (Menini Neto *et al.* 2009; 2016; Leitman *et al.* 2015).

Variables such moisture, light availability, temperature, and seasonality have direct influence in the distribution of epiphytes in the environment (Benzing 1990). Thus, complex variables that are composed by the first ones, for instance in a wide scale, elevation, latitude, continentality and, in a narrow scale, distance from water bodies, stratification on the phorophyte, and relief, also interfere on the epiphyte community.

Low temperature and frost are pointed as limiting to the richness of vascular epiphytes in different scales (Gentry & Dodson 1987; Krömer *et al.* 2005; Blum *et al.* 2011; Hsu *et al.* 2014), which is corroborated in this study, once we found that minimum annual temperature is one of the important variables regarding the obtained floristic relationships. Elevation is directly related with temperature, atmospheric pressure and cloud cover and indirectly related with moisture, sun hours, wind, geology and seasonality (Körner 2004), that is, adds both positive and negative variables to the development of epiphytes, inclusive showing a variation depending upon the epiphytic group. For instance, Orchidaceae and monilophytes present a relative enhancement in the richness following the elevation, reaching a diversity peak in higher altitudes than found in other groups (Moran 1995; Krömer *et al.* 2005).

Positive correlation between geographic distance and Sørensen similarity index is due to the grouping in a cluster of distant areas composed by MOF (since the areas of 'Planalto Meridional', in the Southern Region, grouped together with those present in the Serra da Mantiqueira, in the Southeastern Region of Brazil). This cluster, albeit with a reduced similarity index, contradicts the rare studies that deal with the floristic relationships of the MOF. For instance, studies concerning the flora of shrubs and trees showed great dissimilarities between the areas of MOF of the Southern and Southeastern regions of Brazil (Jarenkow & Budke 2009; Ribeiro 2011).

During the Middle and Upper Holocene (between 4,320 and 1,000 years before the present) the typical tree species of MOF expanded, especially due to the enhancement of moisture, forming forests along the rivers (Bauermann & Behling 2009). Therefore, the MOF of Serra da Mantiqueira took refuge in patches and became isolated from the southern forests, forming islands among the 'campo de altitude', likely due to the dynamic between the field and forest (Behling & Pillar 2007). This isolation, although sufficient for some recognition of distinct floristic sets, as the case of shrubs and trees, seems too weak for homogenisation of the vascular epiphytic flora of MOF with other surrounding physiognomies.

Regarding the environmental variables, the study of Oliveira-Filho *et al.* (2013) also stressed the shortest distance to the ocean, elevation, and variation in the temperature throughout the year, among others, as important in determining the relationships among the forest physiognomies of the Southern Region of Brazil based on flora of trees.

It is possible that the composition of typical tree species in the MOF and, consequently, the probability of being the main phorophytes contribute to higher frequency and sharing of several epiphytic species, explaining the similarity between the areas of the Southern and Southeastern regions, despite the distance between them. Therefore, species like *A. angustifolia* and *P. lambertii*, which are dominant trees in the studied area (Santana 2016), often emphasised among the species of the prominent importance value index in phytosociological studies conducted on MOF of the Southern and Southeastern regions (Geraldi *et al.* 2005; Seger *et al.* 2005; Ribeiro *et al.* 2007; Araujo *et al.* 2010; Silva *et al.* 2012; Souza *et al.* 2012), can possibly be a determinant for the occurrence of epiphytic species shared by areas with this physiognomy, regardless of geographic distance or environmental variables.

Wilberger *et al.* (2009) and Furtado & Menini Neto (2015a) evaluated the vascular epiphytes on *A. angustifolia* and *P. lambertii*, respectively, showing their importance as support for the epiphytic synusia. However, the lack of studies that correlated the occurrence of epiphytes and respective phorophytes in the MOF, regardless of species, impede deeper conclusions about this subject.

Acknowledgements

We wish to thank Dr. Fátima Salimena for support, the Instituto Estadual de Florestas of Minas Gerais (IEF-MG) for the license and logistic support, the Programa de Pós-graduação em Ecologia of Universidade Federal de Juiz de Fora (PGECOL-UFJF) for the logistic support, all the specialists who contributed with identifications, and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the scientific internship of the first author.

References

- Ab'saber AN. 2003. Os domínios de natureza no Brasil: potencialidades paisagísticas. São Paulo, Ateliê Editorial.
- Aguiar LW, Citadini-Zanette V, Martau L, Backes A. 1981. Composição florística de epífitos vasculares numa área localizada nos municípios de Montenegro e Triunfo, Rio Grande do Sul, Brasil. *Iheringia, Série Botânica* 28: 55-93.
- Alves FE, Menini Neto L. 2014. Vascular epiphytes in a forest of Serra da Mantiqueira and floristic relationships with Atlantic high altitude areas in Minas Gerais. *Brazilian Journal of Botany* 37: 187-196.
- APG IV – The Angiosperm Phylogeny Group. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* 181: 1-20.
- Araujo MM, Chami L, Longhi SJ, Avila AL, Brena DA. 2010. Análise de agrupamento em remanescente de Floresta Ombrófila Mista. *Ciência Florestal* 20: 1-18.
- Backes A. 2009. Distribuição geográfica atual da Floresta com Araucária: condicionamento climático. In: Fonseca CR, Souza AF, Leal-Zanchet AM, Dutra T, Backes A, Ganado G. (eds.) *Floresta com Araucária: Ecologia, conservação e desenvolvimento sustentável*. Ribeirão Preto, Holos Editora. p. 137-148.
- Bauermann SG, Behling H. 2009. Dinâmica paleovegetacional da Floresta com Araucária a partir do final do Pleistoceno: o que mostra a palinologia. In: Fonseca CR, Souza AF, Leal-Zanchet AM, Dutra T, Backes A, Ganado G. (eds.) *Floresta com Araucária: Ecologia, conservação e desenvolvimento sustentável*. Ribeirão Preto, Holos Editora. p. 35-38.
- Behling H, Pillar VD. 2007. Late Quaternary vegetation, biodiversity and fire dynamics on the southern Brazilian highland and their implication for conservation and management of modern Araucaria forest and grassland ecosystems. *Philosophical transactions of the Royal Society of London, Biological Sciences* 362: 243-251.
- Benavides AM, Duque AJ, Duivenvoorden JF, Vasco GA, Callejas R. 2005. A first quantitative census of vascular epiphytes in rain forests of Colombian Amazonia. *Biodiversity and Conservation* 14: 739-758.
- Benzing DH. 1990. *Vascular epiphytes*. New York, Cambridge University Press.
- Bhatt A, Gairola S, Govender Y, Baijnath H, Ramdhani S. 2015. Epiphyte diversity on host trees in an urban environment, eThekweni Municipal Area, South Africa. *New Zealand Journal of Botany* 53: 24-37.
- Blum CT, Roderjan CV, Galvão F. 2011. Composição florística e distribuição altitudinal de epífitas vasculares da Floresta Ombrófila Densa na Serra da Prata, Morretes, Paraná, Brasil. *Biota Neotropica* 11: 141-159.
- Bonnet A, Curcio GR, Lavoranti OJ, Galvão F. 2011. Flora epífita vascular em três unidades vegetacionais do rio Tibagi, Paraná, Brasil. *Rodriguésia* 62: 491-498.
- Bonnet A, Caglioni E, Schmitt JL, *et al.* 2013a. Capítulo 1 - Epífitos vasculares da Floresta Ombrófila Densa de Santa Catarina. In: Vibrans AC, Bonnet A, Caglioni E, Gasper AL, Lingner DV. (eds.) *Inventário Florístico Florestal de Santa Catarina*. Vol. 5. Blumenau, Edifurb. p. 27-71.
- Bonnet A, Caglioni E, Schmitt JL, *et al.* 2013b. Capítulo 2 – Descrições das unidades amostrais dos epífitos vasculares. In: Vibrans AC, Bonnet A, Caglioni E, Gasper AL, Lingner DV. (eds.) *Inventário Florístico Florestal de Santa Catarina* Vol. 5. Blumenau, Edifurb. p. 72-335.
- Borgo M, Silva SM. 2003. Epífitos vasculares em fragmentos de Floresta Ombrófila Mista, Curitiba, Paraná, Brasil. *Revista Brasileira de Botânica* 26: 391-401.
- Borgo M, Petean M, Silva SM. 2002. Epífitos vasculares em um remanescente de floresta estacional semidecidual, município de Fênix, PR, Brasil. *Acta Biologica Leopoldensia* 24: 121-130.
- Breier TB. 2005. O epifitismo vascular em florestas do Sudeste do Brasil. Doctorate Thesis, Universidade Estadual de Campinas, Campinas.
- Buzatto CR, Severo BMA, Waechter JL. 2008. Composição florística e distribuição ecológica de epífitos vasculares na Floresta Nacional de Passo Fundo, Rio Grande do Sul. *Iheringia, série Botânica* 63: 231-239.

Vascular epiphytic flora of a high montane environment of Brazilian Atlantic Forest: composition and floristic relationships with other ombrophilous forests

- Cardelús CL, Colwell RK, Watkins Jr. JE. 2006. Vascular epiphyte distribution patterns: explaining the mid-elevation peak richness. *Journal of Ecology* 94: 144-156.
- Carvalho-Silva M. 2008. *Peperomia* Ruiz & Pav. no Brasil: morfologia e taxonomia do subgênero *Rhynchophorum* (Miq.) Dahlst. PhD Thesis, Escola Nacional de Botânica Tropical, Brazil.
- Catchpole D. 2004. The ecology of vascular epiphytes on a *Ficus* L. host (Moraceae) in a Peruvian cloud forest. PhD Thesis, University of Tasmania, Australia.
- Cervi AC, Dombrowski LTD. 1985. Bromeliaceae de um capão de floresta primária do Centro Politécnico de Curitiba (Paraná, Brasil). *Fontqueria* 9: 9-11.
- Cervi AC, Acra LA, Rodrigues L, Train S, Ivanchechen SL, Moreira ALOR. 1988. Contribuição ao conhecimento das epífitas (exclusive Bromeliaceae) de uma floresta de araucária do primeiro planalto paranaense. *Ínsula* 18: 75-82.
- Christenhusz MJM, Zhang XC, Schneider H. 2011. A linear sequence of extant families and genera of lycophytes and ferns. *Phytotaxa* 19: 7-54.
- Dislich R, Mantovani W. 1998. A flora de epífitas vasculares a reserva da Cidade Universitária "Armando de Salles Oliveira" (São Paulo, Brasil). *Boletim de Botânica da Universidade de São Paulo* 17: 1-83.
- Dittrich VAO, Kozena C, Silva SM. 1999. Levantamento florístico de epífitos vasculares no Parque Barigüi, Paraná, Brasil. *Iheringia, série Botânica* 52: 11-22.
- Drummond GM, Martins CS, Machado ABM, Sebaio FA, Antonini Y. 2005. Biodiversidade em Minas Gerais, um atlas para sua conservação. 2nd. edn. Belo Horizonte, Fundação Biodiversitas.
- Felfili JM, Roitman I, Medeiros MM, Sanchez M. 2011. Procedimentos e métodos de amostragem de vegetação. In: Felfili JM, Eisenlohr PV, Melo MMRF, Andrade LA, Meira Neto JAA. (eds.) *Fitossociologia no Brasil: métodos e estudos de casos*. Viçosa, Editora UFV. p. 86-121.
- Filgueiras TS, Nogueira PE, Brochado AL, Guala GF. 1994. Caminhamento: um método expediente para levantamentos florísticos qualitativos. *Cadernos de Geociências* 12: 39-43.
- Forzza RC, Baumgratz JFA, Bicudo CEM, et al. 2012. New Brazilian floristic list highlights conservation challenges. *BioScience* 62: 39-45.
- Freitas L, Salino A, Menini Neto L, et al. 2016. A comprehensive checklist of vascular epiphytes of the Atlantic Forest reveals outstanding endemic rates. *PhytoKeys* 58: 65-79.
- Furtado SG. 2016. Ecologia de epífitas vasculares nas florestas nebulares do Parque Estadual do Ibitipoca, Minas Gerais, Brasil. MSc Thesis, Universidade Federal de Juiz de Fora, Brazil.
- Furtado SG, Menini Neto L. 2013. Epífitas vasculares da floresta ombrófila mista no Parque Estadual da Serra do Papagaio, Minas Gerais, Brasil. Rapid Color Guides Field Museum of Chicago http://fieldguides.fieldmuseum.org/sites/default/files/rapid-color-guides-pdfs/495_2.pdf. 5 Oct. 2013.
- Furtado SG, Menini Neto L. 2015a. Diversity of vascular epiphytes in two high altitude biotopes of the Brazilian Atlantic Forest. *Brazilian Journal of Botany* 38: 295-310.
- Furtado SG, Menini Neto L. 2015b. Diversity of vascular epiphytes in urban environment: a case study in a biodiversity hotspot, the Brazilian Atlantic Forest. *CES Revista* 29: 82-101.
- Gentry AH, Dodson CH. 1987. Diversity and biogeography of neotropical vascular epiphytes. *Annals of the Missouri Botanical Garden* 74: 205-223.
- Geraldi SE, Koehler AB, Kauano EE. 2005. Levantamento fitossociológico de dois fragmentos da Floresta Ombrófila Mista, em Tijucas do Sul, PR. *Revista Acadêmica* 3: 27-36.
- Giongo C, Waechter JL. 2004. Composição florística e estrutura comunitária de epífitos vasculares em uma floresta de galeria na Depressão Central do Rio Grande do Sul. *Revista Brasileira de Botânica* 27: 563-572.
- Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: paleontological Statistics software package for education and data analysis. *Paleontologia Electronica* 4: 1-9.
- Hefler SM, Faustioni P. 2004. Levantamento florístico de epífitos vasculares do Bosque São Cristóvão, Curitiba, Paraná, Brasil. *Revista Estudos de Biologia* 26: 11-19.
- Hertel RJG. 1950. Contribuição à ecologia de flora epífita da serra do mar (vertente oeste) do Paraná. *Arquivos do Museu Paranaense* 8: 3-63.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.
- Hijmans RJ, Guarino L, Cruz M, Rojas E. 2001. Computer tools for spatial analysis of plant genetic resources data: 1. DIVA-GIS. *Plant Genetic Resources Newsletter* 127: 15-19.
- Hobbs RJ, Huenneke LF. 1992. Disturbance, diversity, and invasion: implications for conservation. *Conservation Biology* 6: 324-337.
- Holzner W, Numata M. 1982. Biology and ecology of weeds. The Hague, Dr. W. Junk Publishers.
- Hsu RCC, Wolf JH, Tamis WL. 2014. Regional and elevational patterns in vascular epiphyte richness on an East Asian island. *Biotropica* 46: 549-555.
- Jarenkow JA, Budke JC. 2009. Padrões florísticos e análise estrutural de remanescentes de floresta com Araucária no Brasil. In: Fonseca CR, Souza AF, Leal-Zanchet AM, Dutra TL, Backes A, Ganade G. (eds.) *Floresta com Araucária: ecologia, conservação e desenvolvimento sustentável*. Ribeirão Preto, Holos. p. 35-38.
- Kersten RA. 2006. Epifitismo vascular na bacia do alto Iguaçu, Paraná. PhD Thesis, Universidade Federal do Paraná, Brazil.
- Kersten RA. 2010. Epífitas vasculares – História, participação taxonômica e aspectos relevantes com ênfase na Mata Atlântica. *Hoehnea* 37: 9-38.
- Kersten RA, Kuniyoshi YS, Roderjan CV. 2009. Comunidade epífita em duas formações florestais do Rio São Jerônimo, Bacia do Rio Iguaçu, municípios de Guarapuava e Pinhão, Paraná. *Iheringia, série Botânica* 64: 33-43.
- Kersten RA, Silva SM. 2002. Florística e estrutura do componente epífítico vascular em Floresta Ombrófila Mista Aluvial do rio Barigui, Paraná, Brasil. *Revista Brasileira de Botânica* 25: 259-267.
- Körner C. 2004. Mountain biodiversity, its causes and function. *Ambio* 33: 11-17.
- Krömer T, Kessler M, Gradstein SR, Acebey A. 2005. Diversity patterns of vascular epiphytes along an elevational gradient in the Andes. *Journal of Biogeography* 32: 1799-1809.
- Leitman P, Amorim A, Menini Neto L, Forzza RC. 2014. Epiphytic angiosperms in a mountain forest in southern Bahia, Brazil. *Biota Neotropica* 14: 1-12.
- Leitman P, Amorim A, Sansevero JBB, Forzza RC. 2015. Floristic patterns of epiphytes in the Brazilian Atlantic Forest, a biodiversity hotspot. *Botanical Journal of the Linnean Society* 179: 587-601.
- Lino CF, Albuquerque JL, Dias H. 2007. Cadernos da Reserva da Biosfera da Mata Atlântica nº 32 – Mosaicos de unidade de conservação no corredor da Serra do Mar. São Paulo. São Paulo, Conselho Nacional da Reserva da Biosfera da Mata Atlântica.
- Luer CA. 1986. *Icones Pleurothallidinarum III. Systematics of Pleurothallis (Orchidaceae)*. Monographs in Systematic Botany from the Missouri Botanical Garden, 20. St. Louis, MGB Press.
- Madison M. 1977. Vascular epiphytes: their systematic occurrence and salient features. *Selbyana* 2: 1-13.
- Mania F. 2013. Composição florística de comunidades epífiticas vasculares em Unidades de Conservação no Estado de São Paulo. PhD Thesis, Universidade Estadual Paulista Júlio de Mesquita Filho, Brazil.
- Martinelli G. 2007. Mountain biodiversity in Brazil. *Revista Brasileira de Botânica* 30: 587-597.
- Menini Neto L, Forzza RC, Zappi D. 2009. Angiosperm epiphytes as conservation indicators in forest fragments: A case study from southeastern Minas Gerais, Brazil. *Biodiversity and Conservation* 18: 3785-3807.
- Menini Neto L, Furtado SG, Zappi DC, Oliveira-Filho AT, Forzza RC. 2016. Biogeography of epiphytic Angiosperms in the Brazilian Atlantic Forest, a world biodiversity hotspot. *Brazilian Journal of Botany* 39: 261-173.
- Mittermeier RA, Gil PR, Hoffmann M, et al. 2004. Hotspots revisited: earth's biologically richest and most endangered terrestrial ecoregions. Washington, Ceme.
- Monaco TJ, Weller SC, Ashton FM. 2002. Weed science: principles & practices. New York, John Wiley & Sons, Inc.
- Moran RC. 1995. The importance of mountains to pteridophytes, with emphasis on Neotropical Montane Forests. In: Churchill SP, Balsev H, Forero E, Luteyn JL. (eds.) *Biodiversity and conservation of*



- Neotropical Montane Forests. New York, The New York Botanical Garden. p. 359-363.
- Nadkarni N. 1984. Epiphyte biomass and nutrient capital of a neotropical elfin forest. *Biotropica* 16: 249-256.
- Nieder J, Engwald S, Klawun M, Barthlott W. 2000. Spatial distribution of vascular epiphytes (including hemiepiphytes) in a Lowland Amazon Rain Forest (Surumoni Crane Plot) of Southern Venezuela. *Biotropica* 32: 385-396.
- Oliveira-Filho AT, Budke JC, Jarenkow JA, Eisenlohr PV, Neves DRM. 2013. Delving into the variations in tree species composition and richness across South American subtropical Atlantic and Pampean forests. *Journal of Plant Ecology* 6: 1-23.
- Palmer MW. 1993. Putting things in even better order: the advantages of canonical correspondence analysis. *Ecology* 74: 2215-2230.
- Pereira LC, Chautems A, Mello RM, Menini Neto L. 2013. Gesneriaceae no Parque Estadual da Serra do Papagaio, Minas Gerais, Brasil. *Boletim de Botânica da Universidade de São Paulo* 31: 1-12.
- Petean MP. 2002. Florística e estrutura dos epífitos vasculares em uma área de Floresta Ombrófila Densa Altomontana no Parque Estadual do Pico do Marumbi, Morretes, Paraná, Brasil. MSc Thesis, Universidade Federal do Paraná, Brazil.
- Petean MP. 2009. O componente epífítico vascular em Floresta Ombrófila Densa no litoral paranaense: análise florística, estrutural e de biomassa. PhD Thesis, Universidade Federal do Paraná, Brazil.
- Pridgeon AM. 1982. Numerical analyses in the classification of the Pleurothallidinae (Orchidaceae). *Botanical Journal of the Linnean Society* 85: 103-131.
- Rapini A, Andrade MJG, Giulietti AM, Queiroz LP, Silva JMC. 2009. Introdução. In: Giulietti AM, Rapini A, Andrade MJG, Queiroz LP, Silva JMC (eds.) *Plantas Raras do Brasil*. Belo Horizonte, Conservação Internacional. p. 22-35.
- Ribeiro BR, Longhi SJ, Brena DA, Nascimento ART. 2007. Diversidade e classificação da comunidade arbórea da floresta ombrófila mista da FLONA de São Francisco de Paula, RS. *Ciência Florestal* 17: 101-108.
- Ribeiro TM. 2011. Florística e estrutura da comunidade arbustivo-arbórea em florestas naturais e restauradas com *Araucaria angustifolia* (Bertol.) O. Kuntze no estado de São Paulo, Brasil. MSc Thesis, Universidade Federal de Viçosa, Brazil.
- Rizzini CT. 1997. Tratado de fitogeografia do Brasil. 2nd. edn. Rio de Janeiro, Âmbito Cultural Edições Ltda.
- Rogalski JM, Zanin EM. 2003. Composição florística de epífitos vasculares no estreito de Augusto César, floresta Estacional Decidual do Rio Uruguay, RS, Brasil. *Revista Brasileira de Botânica* 26: 551-556.
- Safford HD. 1999. Brazilian Páramos I. An introduction to the physical environment and vegetation of the *campos de altitude*. *Journal of Biogeography* 26: 693-712.
- Santana LD. 2016. Impacto do incêndio florestal na comunidade arbórea de uma floresta ombrófila mista aluvial altomontana na Serra da Mantiqueira meridional (Minas Gerais). MsC Thesis, Universidade Federal de Juiz de Fora, Brazil.
- Santiago DS. 2013. Composição florística, similaridade e influência de variáveis ambientais de uma floresta de araucária na Serra da Mantiqueira, Minas Gerais, Brasil. MsC Thesis, Universidade Federal de Juiz de Fora, Brazil.
- Saout SL, Hoffmann M, Shi Y, et al. 2013. Protected areas and effective biodiversity conservation. *Science* 342: 803-805.
- Scolforo JR, Mello JM, Silva CPC. 2008. Inventário florestal de Minas Gerais: Floresta Estacional Semidecidual e Ombrófila. Lavras, Editora UFLA.
- Seger CD, Dlugosz FL, Kurasz G, et al. 2005. Levantamento florístico e análise fitossociológica de um remanescente de Floresta Ombrófila Mista localizado no município de Pinhais, Paraná – Brasil. *Floresta* 35: 291-302.
- Silva AC, Higuchi P, Aguiar MD, Negrini M, Fert Neto J, Hess AF. 2012. Relações florísticas e fitossociologia de uma Floresta Ombrófila Mista montana secundária em Lages, Santa Catarina. *Ciência Florestal* 22: 193-206.
- Silva LVC, Viana PL, Mota NFO. 2008. Plano de Manejo do Parque Estadual da Serra do Papagaio, Minas Gerais, Brasil. Belo Horizonte, Instituto Estadual de Florestas.
- Souza FS, Salino A, Viana PL, Salimena FRG. 2012. Pteridófitas da Serra Negra, Minas Gerais, Brasil. *Acta Botanica Brasiliensis* 26: 378-390.
- Stehmann JR, Forzza RC, Salino A, Sobral M, Costa DP, Kamino LHY. 2009. Diversidade taxonômica na Floresta Atlântica. In: Stehmann JR, Forzza RC, Salino A, Sobral M, Costa DP, Kamino LHY. (eds.) *Plantas da Floresta Atlântica*. Rio de Janeiro, Instituto de Pesquisas Jardim Botânico do Rio de Janeiro. p. 3-12.
- ter Braak CJF. 1986. Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67: 1167-1179.
- ter Braak CJF, Smilauer P. 2002. CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). Itaca, Microcomputer Power.
- Thiers B. 2015. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <<http://sweetgum.nybg.org/ih/>> 2 Feb. 2015.
- Watkins Jr. JE, Cardelús C, Colwell RK, Moran RC. 2006. Species richness and distribution of ferns along an elevational gradient in Costa Rica. *American Journal of Botany* 93: 73-83.
- Wilberger TP, Boeni BO, Azambuja CP, et al. 2009. Epífitos vasculares associados à *Araucaria angustifolia*. In: Fonseca CR, Souza AF, Leal-Zanchet AM, Dutra T, Backes A, Ganado G. (eds.) *Floresta com Araucária: Ecologia, conservação e desenvolvimento sustentável*. Ribeirão Preto, Holos Editora. p. 137-147.
- Zotz G. 2013. The systematic distribution of vascular epiphytes – a critical update. *Botanical Journal of the Linnaean Society* 171: 453-481.